


Origination Date: 10/03	Revision Date(s): 10/04, 10/05, 11/06, 11/07, 11/08, 7/10, 7/11
Developed By: Medical Criteria Committee	



Approved:

Robert Mills, MD

Date: 07/28/11

Description:

Percutaneous vertebroplasty is a therapeutic, interventional radiologic procedure performed under imaging guidance that consists of the injection of medical grade cement through a needle into a painful fractured cervical, thoracic or lumbar vertebral body to stabilize the fracture. Vertebroplasty is performed in an attempt to relieve pain and strengthen the spine.

Percutaneous kyphoplasty is similar to vertebroplasty in that stabilization of a collapsed vertebra is accomplished by the injection of bone cement. Under fluoroscopic guidance, an inflatable balloon is inserted to expand a collapsed vertebral body to its natural height prior to the injection of the cement. With kyphoplasty, some of the bony deformity and resulting kyphosis may be reduced which will often significantly improve a patient's pain.

Criteria:

I. Vertebroplasty or kyphoplasty will be covered to plan limitations for patients with vertebral collapse and persistent, debilitating pain in the cervical, thoracic or lumbar bodies resulting from **ANY** of the following:

1. Painful osteoporotic vertebral collapse/compression fractures
2. Traumatic fracture
3. Painful osteolytic vertebral compression fracture related to benign or malignant tumor, such as hemangioma, metastatic disease, myeloma, lymphoma and histiocytosis
4. Steroid induced fractures

AND

II. **ALL** of the following criteria have been met:

1. Severe debilitating pain or loss of mobility that cannot be relieved by at least 6 weeks of optimal medical therapy such as:
 - a. NSAIDS
 - b. Narcotics
 - c. Back bracing
 - d. Physical Therapy
 - e. Initial bed rest with progressive activity; and
2. Other causes of pain, such as herniated intervertebral disk, have been ruled out by CT or MRI; and
3. The affected vertebra has not been extensively destroyed and is at least one-third of its original height; and
4. The vertebral fracture is less than 1 year old

III. Contraindications:

1. Coagulation disorders
2. Underlying infection such as osteomyelitis of the affected vertebra
3. Neurological symptoms related to spinal compression
4. Lack of neurosurgical backup for emergency decompression in the event a neurological deficit develops during the injections of the cement
5. Absence of a confirmed acute or subacute fracture
6. Unstable fracture
7. Asymptomatic vertebral compression fracture

Origination Date: 10/03	Revision Date(s): 10/04, 10/05, 11/06, 11/07, 11/08, 7/10, 7/11
Developed By: Medical Criteria Committee	

8. Burst fracture with retropulsed fragments
9. Known allergy to materials used in either procedure

IV. Information to be Submitted with Pre-Authorization Request:

1. Medical records from the treating physician documenting the spinal level involved, the severity of pain, previous treatments tried, and the patient's neurologic condition
2. X-Ray, CT, or MRI report documenting vertebral collapse

Applicable CPT/HCPC Codes:

Note: list is not all inclusive

22520	Percutaneous vertebroplasty, 1 vertebral body, unilateral or bilateral injection; thoracic
22521	Percutaneous vertebroplasty, 1 vertebral body, unilateral or bilateral injection; lumbar
22522	Percutaneous vertebroplasty, 1 vertebral body, unilateral or bilateral injection; each additional thoracic or lumbar vertebral body (List separately in addition to code for primary procedure)
22523	Percutaneous vertebral augmentation, including cavity creation (fracture reduction and bone biopsy included when performed) using mechanical device, one vertebral body, unilateral or bilateral cannulation (e.g., kyphoplasty); thoracic
22524	Percutaneous vertebral augmentation, including cavity creation (fracture reduction and bone biopsy included when performed) using mechanical device, one vertebral body, unilateral or bilateral cannulation (eg, kyphoplasty); lumbar
22525	Percutaneous vertebral augmentation, including cavity creation (fracture reduction and bone biopsy included when performed) using mechanical device, one vertebral body, unilateral or bilateral cannulation (eg, kyphoplasty); each additional thoracic or lumbar vertebral body (List separately in addition to code for primary procedure)
72291	Radiological supervision and interpretation, percutaneous vertebroplasty, vertebral augmentation, or sacral augmentation (sacroplasty), including cavity creation, per vertebral body or sacrum; under fluoroscopic guidance
72292	Radiological supervision and interpretation, percutaneous vertebroplasty, vertebral augmentation, or sacral augmentation (sacroplasty), including cavity creation, per vertebral body or sacrum; under CT guidance
S2360	Percutaneous vertebroplasty, one vertebral body, unilateral or bilateral injection; cervical
S2361	Each additional cervical vertebral body

References:

- Babb A, Carlson WO. Vertebral compression fractures: treatment and evaluation. S D Med. 2006 Aug;59(8):343-5, 347.
- Balloons for bones. Supplied by the office of Dr. Andrew J. Kokkino
- Bono CM, Heggeness M, Mick C, et al.; North American Spine Society. Newly released

Origination Date: 10/03	Revision Date(s): 10/04, 10/05, 11/06, 11/07, 11/08, 7/10, 7/11
Developed By: Medical Criteria Committee	

vertebroplasty randomized controlled trials: A tale of two trials. Spine J. 2009 Oct 10. [Epub ahead of print]

- Buchbinder R, Osborne RH, Ebeling PR, Wark JD, Mitchell P, Wriedt C, et al. randomized trial of vertebroplasty for painful osteoporotic vertebral fractures. N Engl J Med. 2009 Aug 6;361(6):557-68.
- Burton AW, Mendel E. Vertebroplasty and kyphoplasty. Pain Physicians. 2003 Jul;6(3):335-41. chronic pain, and activity levels. J. Neurosurgery: Spine. Jan. 2003; 98: 36-42.
- Deen HG. Current status of percutaneous vertebral augmentation techniques for vertebral compression fractures. Neurol Neurochir Pol. 2006 July-August; 40(4):320-326.
- Eck JC, Nachtigall D, Humphreys SC, Hodges SD. Comparison of vertebroplasty and balloon kyphoplasty for treatment of vertebral compression fractures: a meta-analysis of the literature. Spine J. 2008 May-Jun;8(3):488-97. Epub 2007 May 29.
- Fourney DR, Schomer DF, Nader R, et al. Percutaneous vertebroplasty and kyphoplasty for painful
- Garfin SR, Buckley RA, Ledlie J, et al. Balloon kyphoplasty for symptomatic vertebral body compression fractures results in rapid, significant, and sustained improvements in back pain, function, and quality of life for elderly patients. Spine 2006 Sep 1;31(19):2213-20.
- Hiwatashi A, Sidhu R, Lee R, et al. Kyphoplasty versus vertebroplasty to increase vertebral body height: a cadaveric study. Radiology. 2005 Dec; 237:1115-1119.
- Hulme PA, Krebs J, Ferguson SJ, et al. Vertebroplasty and kyphoplasty: a systematic review of 69 clinical studies. Spine. 2006 Aug 1;31(17):1983-2001.
- Jensen ME, McGraw Jk, Cardella Jf, Hirsch JA. Position statement on percutaneous vertebral augmentation: a consensus statement developed by the American Society of Interventional and Therapeutic Neuroradiology, Society of Interventional Radiology, American Association of Neurological Surgeons/Congress of Neurological Surgeons, and American Society of Spine Radiology. Vasc Interv Radiol. 2009 Jul;20(7 Suppl):S326-31.
- Karliner L. Balloon kyphoplasty as a treatment for vertebral compression fractures. Technology Assessment. San Francisco, CA: California Technology Assessment Forum (CTAF); June 17, 2009.
- Kasperk C, Grafe IA, Schmitt S, Nöldge G, Weiss C, Da Fonseca K, et al. Three-year outcomes after kyphoplasty in patients with osteoporosis with painful vertebral fractures. J Vasc Interv Radiol. 2010 May;21(5):701-9. Epub 2010 Mar 20.
- Klazen CA, Lohle PN, de Vries J, Jansen FH, Tielbeek AV, Blonk MC, et al. Vertebroplasty versus conservative treatment in acute osteoporotic vertebral compression fractures (Vertos II): an open-label randomised trial. Lancet. 2010 Sep 25;376(9746):1085-92. Epub 2010 Aug 9
- Lane JM, Girardi F, Parvataneni H, et al. Preliminary outcomes of the first 226 consecutive kyphoplasties for the fixation of painful osteoporotic vertebral compression fractures. June 2000. Accessed on the National Osteoporosis Foundation website. Accessed on July 21, 2011 at: www.nof.org
- Ledlie JT, Renfro, M. Balloon kyphoplasty: one-year outcomes in vertebral body height restoration,
- McGuire R. AAOS Clinical Practice Guideline: The Treatment of Symptomatic Osteoporotic Spinal Compression Fractures. J Am Acad Orthop Surg. 2011 Mar;19(3):183-4. Accessed July 21, 2011 at: <http://www.aaos.org/Research/guidelines/guide.asp>
- National Institute for Clinical Excellence (NICE). Balloon kyphoplasty for vertebral compression fractures. Interventional procedure guidance 166. London, UK: NICE; 2006 Apr. Revised 2008 Jan. Accessed Mar 9 2010. Available at URL address: <http://www.nice.org.uk/Guidance/IPG166/Guidance/pdf/English>
- National Institute for Clinical Excellence (NICE). Percutaneous vertebroplasty. Interventional procedure guidance 12. London, UK: NICE; 2003 Nov. Accessed Mar 9 2010. Available at URL address: <http://www.nice.org.uk/page.aspx?o=85733>
- Rousing R, Andersen MO, Jespersen SM, et al. Percutaneous vertebroplasty compared to conservative treatment in patients with painful acute or subacute osteoporotic vertebral fractures: Three-months follow-up in a clinical randomized study. Spine. 2009;34(13):1349-1354.

Origination Date: 10/03	Revision Date(s): 10/04, 10/05, 11/06, 11/07, 11/08, 7/10, 7/11
Developed By: Medical Criteria Committee	

- Society of Interventional Radiology (SIR). Society of Interventional Radiology commentary on vertebroplasty and the August studies in the New England Journal of Medicine. SIR Commentary. Fairfax, VA: SIR; November 24, 2009.
- U.S. Food and Drug Administration (FDA), Center for Devices and Radiological Health (CDRH). Complications related to the use of bone cement and bone void fillers in treating compression fractures of the spine. FDA Public Health Web Notification. Rockville, MD: FDA; updated May 7, 2004. Assessed on July 21, 2011 at: <http://www.fda.gov/cdrh/safety/bonecement.html>. Centers for Medicare and Medicaid Services (CMS). Percutaneous kyphoplasty for vertebral fractures caused by osteoporosis and malignancy. Draft Technology Assessment. Medicare Coverage Database. Baltimore, MD: CMS; 2005. Assessed July 21, 2011 at: <http://www.cms.hhs.gov/mcd/viewtechassess.asp?where=index&tid=25>.
- vertebral body fractures in cancer patients. J. Neurosurgery: Spine. Jan. 2003; 98: 21-30.
- Vertebroplasty benefits selected patients. Hayes Alert; Feb. 2003; VI(2).
- Wardlaw D, Cummings SR, Van Meirhaeghe J, Bastian L, Tillman JB, Ransam J, et al. Efficacy and safety of balloon kyphoplasty compared with non-surgical care for vertebral compression fracture (FREE): a randomised controlled trial. Lancet. 2009 Mar 21;373(9668):1016-24. Epub 2009 Feb 24.
- Physician Advisors