

Origination Date: 8/07	Revision Date(s): 8/08, 9/08
Developed By: Medical Criteria Committee	

Csaba Mera M.D.

Approved: Csaba Mera, MD Date: 09/19/08

Description:

Low back pain is the most frequent cause of disability among individuals under the age of 45 and affects most people at some point in their lives. Conservative management is the first line of treatment for back pain and typically consists of rest, exercise, physical therapy, pain medication, local injections and lumbar bracing. Typically conservative therapy is not recommended in the presence of progressive neurological deficits, when a spinal fracture or dislocation is unstable or for progressive spinal deformity. When conservative management is not successful, surgery may be warranted.

Lumbar fusion was initially used as a treatment for infectious conditions of the spine as well as progressive spinal deformities and traumatic injuries. Other conditions that lumbar fusion has been used to treat are instability as a result of surgical removal of osseous and/or ligamentous structures in the surgical approach for various pathologies, spondylolisthesis, progressive scoliosis, pseudoarthrosis, degenerative disc disease and facet arthropathy.

In performing a lumbar fusion there are different surgical approaches and a variety of implant instrumentation. Bone graft is used to accomplish arthrodesis between lamina, transverse processes, and the vertebral bodies. The grafts can be taken from the patient (autograft), from a donor (allograft) or derived sources (i.e. bone morphogenic proteins or other substances which are used to augment the graft material and have osteo-inductive and/or osteo-conductive properties). Laminar hooks, pedicle screws, interbody devices and other instrumentation are used as an adjunct to fusion to stabilize the spine during the healing of the fusion and to align the spine in a corrected position in the sagittal and frontal planes.

Surgical approaches for lumbar fusion include the following:

From the front and side: Anterior lumbar interbody fusion (ALIF) and eXtreme lateral lumbar interbody fusion (XLIF).

From the back: Posterior lumbar interbody fusion (PLIF), posterior lumbar laminar and inter-transverse process fusion (PLF) or transforaminal lumbar interbody fusion (TLIF).

From the front and back: Combined anterior/posterior lumbar fusion utilizing two approaches.

Minimally invasive: Some surgeons use endoscope to perform minimally invasive decompressive laminotomies, discectomies and lumbar fusion.

Some newer technologies are being introduced which avoid fusion and preserve spinal motion. Such procedures include artificial disc replacement (e.g. Charite and ProDisc), interspinous process decompression systems (e.g. X-STOP) and dynamic stabilization (e.g. Dynesys Spinal System).

Criteria:

ODS will cover lumbar fusion for the following indications:

- I. ODS will cover lumbar fusion for the following conditions when there is associated spinal instability:
 - A. Acute spinal fracture or progressive neurological impairment with instability
 - B. Neural compression after spinal fracture
 - C. Epidural compression or vertebral destruction from a tumor

Origination Date: 8/07	Revision Date(s): 8/08, 9/08
Developed By: Medical Criteria Committee	

- D. Spinal tuberculosis, debridement of the spine for infection, or acute or chronic infection which results in spinal instability
 - E. Pseudoarthrosis (e.g. nonunion of prior fusion)
 - F. Spinal deformity (e.g. congenital, idiopathic, paralytic or disease associated scoliosis and/or kyphosis)
 - G. Instability secondary to intraoperative excessive facet removal, associated radical discectomy, removal of pars interarticularis or pars fracture
 - H. Spondylolisthesis which is symptomatic and interferes with the desired activities of daily living, quality of life, and/or is associated with a manifest or progressive neurologic defect.
- II. **For members who have not had prior lumbar surgery** and have chronic low back pain, associated neurological findings and documented radiological (x-ray, CT, and/or MRI) findings, ODS will cover lumbar fusions when **all** of the following criteria are met:
- A. The member has tried at least 3 months of conservative therapy including rest, pain medications, physical therapy, and local injections; **and**
 - B. The surgeon requesting the lumbar fusion has evaluated the patient on at least two occasions prior to requesting the fusion; **and**
- **Note: if the patient has a progressive neurological deficit, both criteria above can be waived.
- C. Mechanical (non-radicular) low back pain with instability as demonstrated on flexion/extension and/or side bending x-rays; **or**
 - D. Spondylolisthesis which is symptomatic and interferes with the desired activities of daily living, quality of life, and/or is associated with a manifest or progressive neurological deficit. This would include congenital, isthmic, degenerative, traumatic, and neoplastic spondylolisthesis; **or**
 - E. The member has chronic, severe and significantly disabling low back pain from degenerative disc disease which has failed 6 months of conservative treatment and has documented radiological (e.g. x-ray, CT, and/or MRI) evidence of disease.
- III. **For members who have had prior laminectomy, discectomy, or other decompressive procedure at the same level**, ODS will cover lumbar when the following criteria is met:
- A. 3 months of conservative therapy has failed to relieve symptoms; **and**
 - B. Mechanical (non-radicular) low back pain with instability (as defined above in II C); **or**
 - C. Mechanical (non-radicular) low back pain with spondylolisthesis, sagittal, coronal, and/or rotational deformity, or other condition leading to a progressive, measurable deformity; **or**
 - D. Objective signs/symptoms compatible with neurogenic claudication or lumbar radiculopathy that is supported by MRI or CT and neurological exam; **or**
 - E. Evidence of a post-laminectomy structural deficiency.
- IV. **For members who have had prior fusion at the same level**, ODS will cover lumbar fusion when **one** of the following criteria are met:
- A. Pseudoarthrosis with or without hardware failure; **or**
 - B. Neurogenic symptoms or lumbar radiculopathy supported by MRI, CT or myelography and a detailed clinical neurological examination.
- V. **For member who have had prior fusion at an adjacent level(s)**, ODS will cover lumbar fusion if conservative therapy has failed to relieve symptoms and findings require additional levels to be included in the fusion mass. The member must meet the same criteria as described in II above for patients with no prior history of spinal surgery.

Origination Date: 8/07	Revision Date(s): 8/08, 9/08
Developed By: Medical Criteria Committee	

Limitations:

- For artificial disc replacement requests please see ODS medical necessity criteria for Intervertebral Disc Prosthesis.
- For requests for X-STOP, please see ODS medical necessity criteria for Interspinous Process Decompression System (e.g. X-STOP).

Not Covered:

ODS considers the following experimental and investigational because there is insufficient evidence in peer reviewed medical literature to establish their effectiveness:

- Dynamic spine stabilization device systems (e.g. Dynesys)

Information to be Submitted with Pre-Authorization Request:

- Chart notes from the treating physician including subjective and objective symptoms and findings.
- Radiographic study reports
- Conservative treatment attempts
- Prior history of spine surgery or other treatment

References:

- Agency for Healthcare Research and Quality (AHRQ). Treatment of degenerative lumbar spinal stenosis: summary, evidence report/technology assessment: number 32. AHRQ publication no. 01-E047. Rockville, MD: Agency for Healthcare Research and Quality; 2001 Mar. Accessed July 6, 2007. Available at URL address: <http://www.ahrq.gov/clinical/epcs/sums/stenosum.htm>
- U.S. Food and Drug Administration (FDA). Centerpulse Spin-Tech, Inc. Dynesys Spinal System. 510(k) summary. K031511. March 5, 2004. Accessed July 9, 2007. Available at URL address: <http://www.fda.gov/cdrh/pdf3/k031511.pdf>
- Vinas F. Lumbar spine fractures and dislocations. eMedicine. Updated June 14, 2006. Accessed July 9, 2007. Available at URL address: <http://www.emedicine.com/orthoped/topic176.htm>.
- Frantzides CT, Zeni TM, Phillips FM, et al. L5-S1 laparoscopic anterior interbody fusion. JLS. 2006 Oct-Dec;10(4):488-92
- Anderson PA, Tribus CB, Kitchel SH. Treatment of neurogenic claudication by interspinous decompression: application of the X STOP device in patients with lumbar degenerative spondylolisthesis. J Neurosurg Spine. 2006 Jun; 4(6):463-71.
- Welch WC, Cheng BC, Awad TE, et al. Clinical outcomes of the Dynesys dynamic neutralization system: 1-year preliminary results. Neurosurg Focus. 2007 Dec 15;22(1):E8.
- Moller H, Hedlund R, Surgery versus conservative management in adult isthmic spondylolisthesis-a prospective randomized study: part 1. Spine. 2000 Jul 1;25(13):1711-5.
- Silber JS, Anderson DG, Hayes VM, Vaccaro AR. Advances in surgical management of lumbar degenerative disease. Orthopedics. 2002 Jul; 25(7):767-71.
- Fritzell P, Hagg O, Wessberg P, Nordwall A. Lumbar fusion versus nonsurgical treatment for chronic low back pain: a multicenter randomized controlled trial from the Swedish Lumbar Spine Study Group. Spine. 2001 Dec 1;26(23):2521-32.
- Ivar Brox J, Sorensen R, Friis A, et al. Randomized clinical trial of lumbar instrumented fusion and cognitive intervention and exercises in patients with chronic low back pain and disc degeneration. Spine. 2003 Sep 1;28(17):1913-21.
- Guidelines for lumbar fusion (arthrodesis). National Guideline Clearing house. Accessed on July 9, 2007. Available at URL address: http://www.guidelines.gov/summary/symmary.aspx?doc_id=4210&nbr=003218&sting=lumbar+AND+fusion.
- Deyo R, Nachemson A, Mirza S. Spinal fusion surgery-the case for restraint. N Engl J Med. 2004 Feb 12;350(7):722-726.
- Peer review by a board certified orthopedic surgeon and professor of orthopedics.

Origination Date: 8/07	Revision Date(s): 8/08, 9/08
Developed By: Medical Criteria Committee	

- Physician Advisors